



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$2(x-r)+2(mx-r)+2r=x+mx+x\sqrt{1+m^2}.$$

$$\therefore r=\frac{1}{2}x[1+m-\sqrt{1+m^2}].$$

$$GH=\frac{1}{3}BC=\frac{1}{3}mx, \quad KH=\frac{1}{2}AB-HB=\frac{1}{2}x-r.$$

$$\therefore KH=\frac{1}{2}x[\sqrt{1+m^2}-m], \quad GK=\frac{1}{3}CK=\frac{1}{3}x\sqrt{1+m^2}.$$

$$\therefore KG^2=GH^2+KH^2, \text{ or } \frac{1}{9}x^2(1+m^2)=\frac{1}{9}m^2x^2+\frac{1}{4}x^2[\sqrt{1+m^2}-m]^2.$$

$$\therefore \frac{1}{4}[\sqrt{1+m^2}-m]^2=\frac{1}{36}, \text{ or } \sqrt{1+m^2}-m=\frac{1}{6}. \quad \therefore m=\frac{4}{3}.$$

$$\therefore AB:BC=3:4, \text{ and } HB:AB:BC:AC=1:3:4:5.$$

Also solved by J. C. NAGLE.

### DIOPHANTINE ANALYSIS.

60. Proposed by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics, The Russell College, Lebanon, Va.

It is required to find six positive numbers, such that if each be diminished by five-half times the fifth power of their sum the six remainders will be rational fifth powers.

Solution by the PROPOSER.

Let  $u, v, w, x, y, z$ , be the six numbers required, and let  $u+v+w+x+y+z=s$ .

$$\text{Then } u-\frac{5}{2}s^5=h^5s^5/q^5, \quad v-\frac{5}{2}s^5=k^5s^5/q^5, \quad w-\frac{5}{2}s^5=l^5s^5/q^5,$$

$$x-\frac{5}{2}s^5=m^5s^5/q^5, \quad y-\frac{5}{2}s^5=n^5s^5/q^5, \quad z-\frac{5}{2}s^5=p^5s^5/q^5,$$

Adding these six equations we get

$$s-15s^5=(s^5/q^5)(h^5+k^5+l^5+m^5+n^5+p^5).$$

$$\text{Let } h^5+k^5+l^5+m^5+n^5+p^5=q^5. \quad \therefore s=\frac{1}{2}.$$

$$\therefore u=\frac{1}{32}[\frac{5}{2}+(h^5/q^5)], \quad v=\frac{1}{32}[\frac{5}{2}+(k^5/q^5)], \quad w=\frac{1}{32}[\frac{5}{2}+(l^5/q^5)],$$

$$x=\frac{1}{32}[\frac{5}{2}+(m^5/q^5)], \quad y=\frac{1}{32}[\frac{5}{2}+(n^5/q^5)], \quad z=\frac{1}{32}[\frac{5}{2}+(p^5/q^5)].$$

$$\text{Let } h=4, k=5, l=6, m=7, n=9, p=11, q=12.$$

$$u=\frac{1^2 2^1 7}{15552}, \quad v=\frac{6^2 5^2 0^5}{7962624}, \quad w=\frac{8^1}{1024}, \quad x=\frac{6^3 8^8 8^7}{7962627}, \quad y=\frac{9^2 0^3}{32768}, \quad z=\frac{7^8 3^1 3^1}{7962624}.$$

$$\text{Let } h=5, k=10, l=11, m=16, n=19, p=29, q=30.$$

$$u=\frac{1^9 4^4 1}{248832}, \quad v=\frac{1^2 1^7}{15552}, \quad w=\frac{6^0 9^1 1^0 5^1}{77600000}, \quad x=\frac{3^8 6^2 4^1 1}{48600000}, \quad y=\frac{6^3 2^2 0^0 0^0}{77600000}, \quad z=\frac{8^1 2^6 1^1 4^0}{77760000}.$$

### AVERAGE AND PROBABILITY.

58. Proposed by HENRY HEATON, M. Sc., Atlantic, Iowa.

From a point on the surface of a circle two lines are drawn to the circumference. Required the average area that may be cut from the circle in this way if the lines are supposed to be drawn at equal angular intervals.

Query I. How does this differ from problem 32?

Query II. Is *sector* the proper word to use for the surface thus cut off?

Query III. Is it absolutely correct to use the word *random* in average problems?